arXiv:1704.08183v1 [math.CA] 14 Apr 2017

A Dunkl Analogue of Operators Including Two-variable Hermite polynomials

Rabia Aktaş, Bayram Çekim, and Fatma Taşdelen

Abstract. The aim of this paper is to introduce a Dunkl generalization of the operators including two variable Hermite polynomials which are defined by Krech [**14**] (Krech, G. A note on some positive linear operators associated with the Hermite polynomials, Carpathian J. Math., 32 (1) (2016), 71–77) and to investigate approximating properties for these operators by means of the classical modulus of continuity, second modulus of continuity and Peetre's K-functional.

1. Introduction

Up to now, linear positive operators and their approximation properties have been studied by many research workers, see for example [**4**], [**5**], [**6**], [**9**], [**10**], [**19**], [**22**] and references therein. Also, linear positive operators defined via generating functions and their further extensions are intensively studied by a large number of authors. For various extensions and further properties, we refer for example Altin et.al [**1**], Dogru et. al [**8**], Olgun et.al [**17**], Sucu et. al [**21**], Tasdelen et.al [**23**], Varma et.al [**24, 25**].

Recently, linear positive operators generated by a Dunkl generalization of the exponential function have been stated by many authors. In [**20**], Dunkl analogue of Szász operators by using Dunkl analogue of exponential function was given as follows

$$S_n^*(g;x)=\frac{1}{e_\nu(nx)}\sum_{k=0}^{\infty}\frac{(nx)^k}{\gamma_\nu(k)}g\left(\frac{k+2\nu\theta_k}{n}\right)\ ;\ n\in\mathbb{N},\ \nu,\ x\in[0,\infty) \tag{1.1}$$

for $g\in C[0,\infty)$ where Dunkl analogue of exponential function is defined by

$$e_\nu(x)=\sum_{k=0}^{\infty}\frac{x^k}{\gamma_\nu(k)} \tag{1.2}$$

for $k\in\mathbb{N}_0$ and $\nu>-\frac{1}{2}$ and the coefficients γ_ν are as follows

$$\gamma_\nu(2k)=\frac{2^{2k}k!\Gamma(k+\nu+1/2)}{\Gamma(\nu+1/2)}\text{ and }\gamma_\nu(2k+1)=\frac{2^{2k+1}k!\Gamma(k+\nu+3/2)}{\Gamma(\nu+1/2)} \tag{1.3}$$

2000 *Mathematics Subject Classification.* Primary 41A25, 41A36; Secondary 33C45.

Key words and phrases. Dunkl analogue, Hermite polynomial, modulus of continuity, Korovkin's type approximation theorem.

in [**18**]. Also, the coefficients γ_ν verify the recursion relation

$$\frac{\gamma_\nu(k+1)}{\gamma_\nu(k)} = (2\nu\theta_{k+1} + k + 1), \ k \in \mathbb{N}_0, \tag{1.4}$$

where

$$\theta_k = \begin{cases} 0, & if\ k = 2p \\ 1, & if\ k = 2p+1 \end{cases} \tag{1.5}$$

for $p \in \mathbb{N} \cup \{0\}$. Similarly, Stancu-type generalization of Dunkl analogue of Szász-Kantorovich operators and Dunkl generalization of Szász operators via q-calculus have been defined in [**11, 12**] and for other research see [**15, 16**].

The two-variable Hermite Kampe de Feriet polynomials $H_n(\xi, \alpha)$ are defined by (see [**3**])

$$\sum_{n=0}^{\infty} \frac{H_n(\xi,\alpha)}{n!} t^n = e^{\xi t + \alpha t^2}$$

from which, it follows

$$H_n(\xi,\alpha) = n! \sum_{k=0}^{\left[\frac{n}{2}\right]} \frac{\alpha^k \xi^{n-2k}}{k!(n-2k)!}.$$

In a recent paper, Krech [**14**] has introduced the class of operators G_n^α given by

$$G_n^\alpha(f;x) = e^{-\left(nx+\alpha x^2\right)} \sum_{k=0}^{\infty} \frac{x^k}{k!} H_k(n,\alpha) f\left(\frac{k}{n}\right) \ , \ x \in \mathbb{R}_0^+ \ , \ n \in \mathbb{N} \ \ , \alpha \geq 0 \tag{1.6}$$

in terms of two variable Hermite polynomials and investigated approximation properties of G_n^α .

In the present paper, we first give the Dunkl generalization of two variable Hermite polynomials and then we define a class of operators by using the Dunkl generalization of two variable Hermite polynomials. We give the rates of convergence of the operators T_n to f by means of the classical modulus of continuity, second modulus of continuity and Peetre's K-functional and in terms of the elements of the Lipschitz class $Lip_M(\alpha)$.

2. The Dunkl generalization of two variable Hermite polynomials

The Dunkl generalization of two variable Hermite polynomials is defined by

$$\sum_{n=0}^{\infty} \frac{H_n^\mu(\xi,\alpha)}{n!} t^n = e^{\alpha t^2} e_\mu(\xi t) \tag{2.1}$$

from which, we conclude

$$H_n^\mu(\xi,\alpha) = n! \sum_{k=0}^{\left[\frac{n}{2}\right]} \frac{\alpha^k \xi^{n-2k}}{k! \gamma_\mu(n-2k)},$$

which gives the two variable Hermite polynomials as $\mu = 0$. For our purpose, we denote

$$h_n^\mu(\xi,\alpha) = \frac{\gamma_\mu(n) H_n^\mu(\xi,\alpha)}{n!}$$

and we can write that the polynomials $h_n^{\mu}(\xi,\alpha)$ are generated by

$$\sum_{n=0}^{\infty}\frac{h_n^{\mu}(\xi,\alpha)}{\gamma_{\mu}(n)}t^n=e^{\alpha t^2}e_{\mu}(\xi t) \tag{2.2}$$

where

$$h_n^{\mu}(\xi,\alpha)=\gamma_{\mu}(n)\sum_{k=0}^{\left[\frac{n}{2}\right]}\frac{\alpha^k\xi^{n-2k}}{k!\gamma_{\mu}(n-2k)}.$$

In order to obtain some properties of $h_n^{\mu}(\xi,\alpha)$, we remind the following definition and lemma given in [**18**].

DEFINITION 1. [**18**] *Let $\mu\in\mathbb{C}_0$ ($\mathbb{C}_0:=\mathbb{C}\setminus\left\{-\frac{1}{2},-\frac{3}{2},...\right\}$, $x\in\mathbb{C}$ and let φ be entire function. The linear operator $\mathbb{D}_{\mu}$ is defined on all entire functions φ on $\mathbb{C}$ by*

$$\mathbb{D}_{\mu}(\varphi(x))=\varphi^{'}(x)+\frac{\mu}{x}(\varphi(x)-\varphi(-x)),\ x\in\mathbb{C}. \tag{2.3}$$

We use the notation $\mathbb{D}_{\mu,x}$ since $\mathbb{D}_{\mu}$ is acting on functions of the variable x. Thus, $\mathbb{D}_{\mu,x}(\varphi(x))=(\mathbb{D}_{\mu}\varphi)(x)$.

LEMMA 1. [**18**] *Let φ,ψ be entire functions. For the linear operator $\mathbb{D}_{\mu}$, the following statements hold*

i) $\mathbb{D}_{\mu}^{j}:x^n\to\frac{\gamma_{\mu}(n)}{\gamma_{\mu}(n-j)}x^{n-j},j=0,1,2,...,n\ (n\in\mathbb{N});\ \mathbb{D}_{\mu}^{j}:1\to 0$

ii) $\mathbb{D}_{\mu}(\varphi\psi)=\mathbb{D}_{\mu}(\varphi)\psi+\varphi\mathbb{D}_{\mu}(\psi)$, *where φ is an even function*

iii) $\mathbb{D}_{\mu}:e_{\mu}(\lambda x)\to\lambda e_{\mu}(\lambda x)$.

By using these definition and lemma, we can state the next result.

LEMMA 2. *For the Dunkl generalization of two variable Hermite polynomials $h_n^{\mu}(\xi,\alpha)$, the following results hold true*

(i) $\sum_{n=0}^{\infty}\frac{h_{n+1}^{\mu}(\xi,\alpha)}{\gamma_{\mu}(n)}t^n=(\xi+2\alpha t)e^{\alpha t^2}e_{\mu}(\xi t)$

(ii) $\sum_{n=0}^{\infty}\frac{h_{n+2}^{\mu}(\xi,\alpha)}{\gamma_{\mu}(n)}t^n=(\xi^2+4\xi\alpha t+4\alpha^2t^2+2\alpha)e^{\alpha t^2}e_{\mu}(\xi t)+4\alpha\mu e^{\alpha t^2}e_{\mu}(-\xi t)$

PROOF. Applying the linear operator $\mathbb{D}_{\mu}$ in view of Lemma 1 , we have

$$\begin{array}{l}\mathbb{D}_{\mu}(te_{\mu}(\xi t))=(t\xi+1)e_{\mu}(\xi t)+2\mu e_{\mu}(-\xi t)\\ \mathbb{D}_{\mu}(e^{\alpha t^2})=2\alpha te^{\alpha t^2}.\end{array} \tag{2.4}$$

Also applying the linear operator $\mathbb{D}_{\mu}$ to both side of the generating function (2.2), we have

$$\sum_{n=0}^{\infty}\frac{h_n^{\mu}(\xi,\alpha)}{\gamma_{\mu}(n)}\mathbb{D}_{\mu}(t^n)=\mathbb{D}_{\mu}(e^{\alpha t^2}e_{\mu}(\xi t)).$$

By using (2.4) and Lemma 1 (i), we get the first relation. Similarly, if we apply the linear operator $\mathbb{D}_{\mu}$ to the relation in (i), we get

$$\sum_{n=0}^{\infty}\frac{h_{n+1}^{\mu}(\xi,\alpha)}{\gamma_{\mu}(n)}\mathbb{D}_{\mu}(t^n)=\mathbb{D}_{\mu}\left[(\xi+2\alpha t)e^{\alpha t^2}e_{\mu}(\xi t)\right]$$

from (2.4) and Lemma 1, it follows

$$\sum_{n=0}^{\infty}\frac{h_{n+2}^{\mu}(\xi,\alpha)}{\gamma_{\mu}(n)}t^n=(\xi^2+4\xi\alpha t+4\alpha^2t^2+2\alpha)e^{\alpha t^2}e_{\mu}(\xi t)+4\alpha\mu e^{\alpha t^2}e_{\mu}(-\xi t).$$

□

DEFINITION 2. *With the help of the Dunkl generalization of two variable Hermite polynomials given in (2.2), we introduce the operators* $T_n(f;x)$, $n\in\mathbb{N}$ *given by*

$$T_n(f;x):=\frac{1}{e^{\alpha x^2}e_{\mu}(nx)}\sum_{k=0}^{\infty}\frac{h_k^{\mu}(n,\alpha)}{\gamma_{\mu}(k)}x^k f\left(\frac{k+2\mu\theta_k}{n}\right) \tag{2.5}$$

where $\alpha\geq 0,\mu\geq 0$ *and* $x\in[0,\infty)$. *The operators (2.5) are linear and positive. In the case of* $\mu=0$, *it gives* G_n^{α} *given by (1.6)*

LEMMA 3. *For the operators* $T_n(f;x)$, *we can obtain the following equations:*

(i) $T_n(1;x)=1$
(ii) $T_n(t;x)=x+\frac{2\alpha x^2}{n}$
(iii) $T_n(t^2;x)=x^2+\frac{4\alpha}{n^2}x^2+\frac{4\alpha}{n}x^3+\frac{4\alpha^2}{n^2}x^4+\frac{x}{n}+\frac{2\mu x}{n}\frac{e_{\mu}(-nx)}{e_{\mu}(nx)}$

PROOF. By using the generating function in (2.2), the relation (i) holds. For the proof of (ii), in view of the recursion relation in (1.4), we get

$$T_n(t;x)=\frac{1}{ne^{\alpha x^2}e_{\mu}(nx)}\sum_{k=1}^{\infty}\frac{h_k^{\mu}(n,\alpha)}{\gamma_{\mu}(k-1)}x^k.$$

When we replace k by $k+1$, we obtain (ii) by use of Lemma 2 (i). For the proof of (iii), by using (1.4), we have

$$T_n(t^2;x)=\frac{x}{n^2e^{\alpha x^2}e_{\mu}(nx)}\sum_{k=0}^{\infty}(k+1+2\mu\theta_{k+1})\frac{h_{k+1}^{\mu}(n,\alpha)}{\gamma_{\mu}(k)}x^k.$$

From the equation

$$\theta_{k+1}=\theta_k+(-1)^k, \tag{2.6}$$

it yields

$$\begin{aligned}T_n(t^2;x) &= \frac{x}{n^2e^{\alpha x^2}e_{\mu}(nx)}\sum_{k=0}^{\infty}(k+2\mu\theta_k)\frac{h_{k+1}^{\mu}(n,\alpha)}{\gamma_{\mu}(k)}x^k\\ &\quad+\frac{x}{n^2e^{\alpha x^2}e_{\mu}(nx)}\sum_{k=0}^{\infty}(1+2\mu(-1)^k)\frac{h_{k+1}^{\mu}(n,\alpha)}{\gamma_{\mu}(k)}x^k.\end{aligned}$$

Using the recursion relation in (1.4) in the first series, it follows

$$\begin{aligned}T_n(t^2;x) &= \frac{x^2}{n^2e^{\alpha x^2}e_{\mu}(nx)}\sum_{k=0}^{\infty}\frac{h_{k+2}^{\mu}(n,\alpha)}{\gamma_{\mu}(k)}x^k+\frac{x}{n^2e^{\alpha x^2}e_{\mu}(nx)}\sum_{k=0}^{\infty}\frac{h_{k+1}^{\mu}(n,\alpha)}{\gamma_{\mu}(k)}x^k\\ &\quad+\frac{2\mu x}{n^2e^{\alpha x^2}e_{\mu}(nx)}\sum_{k=0}^{\infty}(-x)^k\frac{h_{k+1}^{\mu}(n,\alpha)}{\gamma_{\mu}(k)}\end{aligned}$$

from Lemma 2 (i) and (ii), we complete the proof of (iii). □

LEMMA 4. *As a consequence of Lemma 3, we can give the next results for T_n operators*

$$\begin{aligned}\Delta_1 &= T_n(t-x;x) = \frac{2\alpha x^2}{n} \\ \Delta_2 &= T_n((t-x)^2;x) = \frac{1}{n^2}x\left(4x^3\alpha^2+4\alpha x+n\right)+\frac{2\mu x}{n}\frac{e_\mu(-nx)}{e_\mu(nx)} \end{aligned} \tag{2.7}$$

THEOREM 1. *For T_n operators and any uniformly continuous bounded function g on the interval $[0,\infty)$, we can give*

$$T_n(g;x) \overset{uniformly}{\rightrightarrows} g(x)$$

on each compact set $A \subset [0,\infty)$ when $n \to \infty$.

PROOF. From Korovkin Theorem in [**13**], when $n \to \infty$, we have $T_n(g;x) \overset{\text{uniformly}}{\rightrightarrows} g(x)$ on $A \subset [0,\infty)$ which is each compact set because $\lim_{n\to\infty} T_n(e_i;x) = x^i$, for $i = 0,1,2$, which is uniformly on $A \subset [0,\infty)$ with the help of using Lemma 4. □

THEOREM 2. *The operator T_n maps $C_B(\mathbb{R}_0^+)$ into $C_B(\mathbb{R}_0^+)$ and $\|T_n(f)\| \leq \|f\|$ for each $f \in C_B(\mathbb{R}_0^+)$.*

3. Convergence of operators in (2.5)

In what follows we give some rates of convergence of the operators T_n. Firstly, we recall some definitions as follows. Let $Lip_M(\alpha)$ Lipschitz class of order α. If $g \in Lip_M(\alpha)$, the inequality

$$|g(s)-g(t)| \leq M|s-t|^\alpha$$

holds where $s,t \in [0,\infty)$, $0 < \alpha \leq 1$ and $M > 0$. $\widetilde{C}[0,\infty)$ is the space of uniformly continuous on $[0,\infty)$. The modulus of continuity $g \in \widetilde{C}[0,\infty)$ is denoted by

$$\omega(g;\delta) := \sup_{\substack{s,t\in[0,\infty)\\ |s-t|\leq\delta}} |g(s)-g(t)|. \tag{3.1}$$

We first estimate the rates of convergence of the operators T_n by using modulus of continuity and in terms of the elements of the Lipschitz class $Lip_M(\alpha)$.

THEOREM 3. *If $h \in Lip_M(\alpha)$, we have*

$$|T_n(h;x)-h(x)| \leq M(\Delta_2)^{\alpha/2}$$

where Δ_2 is given in Lemma 4.

PROOF. Since $h \in Lip_M(\alpha)$, it follows from linearity

$$|T_n(h;x)-h(x)| \leq T_n(|h(t)-h(x)|;x) \leq MT_n(|t-x|^\alpha;x).$$

From Lemma 4 and Hölder's famous inequality, we can write

$$|T_n(h;x)-h(x)| \leq M[\Delta_2]^{\frac{\alpha}{2}}.$$

Thus, we find the required inequality. □

THEOREM 4. *The operators in (2.5) verify the inequality*

$$|T_n(g;x)-g(x)| \leq \left(1+\sqrt{\frac{1}{n}x\left(4x^3\alpha^2+4x\alpha+n\right)+2\mu x\frac{e_\mu(-nx)}{e_\mu(nx)}}\right)\omega\left(g;\frac{1}{\sqrt{n}}\right),$$

where $g \in \widetilde{C}[0,\infty)$.

PROOF. By Lemma 4, from Cauchy-Schwarz's inequality and the property of modulus of continuity

$$|g(t)-g(x)| \leq w(g;\delta)\left(\frac{|t-x|}{\delta}+1\right), \tag{3.2}$$

it follows

$$\begin{aligned} |T_n(g;x)-g(x)| &\leq T_n(|g(t)-g(x)|;x) \\ &\leq \left(1+\frac{1}{\delta}T_n(|t-x|;x)\right)\omega(g;\delta) \\ &\leq \left(1+\frac{1}{\delta}\sqrt{\Delta_2}\right)\omega(g;\delta). \end{aligned}$$

Then from Lemma 4, one has

$$|T_n(g;x)-g(x)| \leq \left(1+\frac{1}{\delta}\sqrt{\frac{1}{n^2}x\left(4x^3\alpha^2+4\alpha x+n\right)+\frac{2\mu x}{n}\frac{e_\mu(-nx)}{e_\mu(nx)}}\right)\omega(g;\delta), \tag{3.3}$$

by choosing $\delta=\frac{1}{\sqrt{n}}$, we completes the proof. □

Let $C_B[0,\infty)$ denote the space of uniformly continuous and bounded functions on $[0,\infty)$. Also

$$C_B^2[0,\infty)=\{g\in C_B[0,\infty): g',g''\in C_B[0,\infty)\} \tag{3.4}$$

with the norm

$$\|g\|_{C_B^2[0,\infty)}=\|g\|_{C_B[0,\infty)}+\|g'\|_{C_B[0,\infty)}+\|g''\|_{C_B[0,\infty)}$$

for $\forall g\in C_B^2[0,\infty)$.

LEMMA 5. *For* $h\in C_B^2[0,\infty)$*, the following inequality holds true*

$$|T_n(h;x)-h(x)| \leq [\Delta_1+\Delta_2]\|h\|_{C_B^2[0,\infty)}, \tag{3.5}$$

where Δ_1 *and* Δ_2 *are given by in Lemma 4.*

PROOF. From the Taylor's series of the function h,

$$h(s)=h(x)+(s-x)h'(x)+\frac{(s-x)^2}{2!}h''(\varrho),\ \varrho\in(x,s).$$

Applying the operator T_n to both sides of this equality and then using the linearity of the operator, we have

$$T_n(h;x)-h(x)=h'(x)\Delta_1+\frac{h''(\varrho)}{2}\Delta_2.$$

From Lemma 4, it yields

$$\begin{aligned}|T_n(h;x)-h(x)| &\leq \frac{2\alpha x^2}{n}\|h'\|_{C_B[0,\infty)} \\ &\quad+\left[\frac{1}{n^2}x\left(4x^3\alpha^2+4\alpha x+n\right)+\frac{2\mu x}{n}\frac{e_\mu(-nx)}{e_\mu(nx)}\right]\|h''\|_{C_B[0,\infty)} \\ &\leq [\Delta_1+\Delta_2]\|h\|_{C_B^2[0,\infty)},\end{aligned}$$

which finishes the proof. □

Now we recall that the second order of modulus continuity of f on $C_B[0,\infty)$ is given as

$$\omega_2(f;\delta):=\sup_{0<s\leq\delta}\|f(.+2s)-2f(.+s)+f(.)\|_{C_B[0,\infty)}.$$

Peetre's K-functional of the function $f\in C_B[0,\infty)$ is as follows

$$K(f;\delta):=\inf_{g\in C_B^2[0,\infty)}\left\{\|f-g\|_{C_B}+\delta\|g\|_{C_B^2}\right\} \tag{3.6}$$

The relation between K and ω_2 is as

$$K(f;\delta)\leq M\left\{w_2\left(f;\sqrt{\delta}\right)+\min(1,\delta)\|f\|_{C_B}\right\} \tag{3.7}$$

for all $\delta>0$. Here M is a positive constant. Now, we can give the important theorem.

THEOREM 5. *For the operators by (2.5), the following inequality holds*

$$|T_n(g;x)-g(x)|\leq 2M\left\{\min\left(1,\frac{\chi_n(x)}{2}\right)\|g\|_{C_B[0,\infty)}+\omega_2\left(g;\sqrt{\frac{\chi_n(x)}{2}}\right)\right\} \tag{3.8}$$

where $\forall g\in C_B[0,\infty)$, $x\in[0,\infty)$, *M is a positive constant which is independent of n and* $\chi_n(x)=\Delta_1+\Delta_2$.

PROOF. For any $f\in C_B^2[0,\infty)$, from the triangle inequality, we can write

$$\Theta=|T_n(g;x)-g(x)|\leq|T_n(g-f;x)|+|T_n(f;x)-f(x)|+|g(x)-f(x)|$$

from Lemma 5, which follows

$$\begin{aligned}\Theta &\leq 2\|g-f\|_{C_B[0,\infty)}+\chi_n(x)\|f\|_{C_B^2[0,\infty)} \\ &= 2\left\{\|g-f\|_{C_B[0,\infty)}+\frac{\chi_n}{2}(x)\|f\|_{C_B^2[0,\infty)}\right\}.\end{aligned}$$

From (3.6), we have

$$\Theta\leq 2K\left(g;\frac{\chi_n(x)}{2}\right),$$

which holds

$$\Theta\leq 2M\left\{\min\left(1,\frac{\chi_n(x)}{2}\right)\|g\|_{C_B[0,\infty)}+\omega_2\left(g;\sqrt{\frac{\chi_n(x)}{2}}\right)\right\}$$

from (3.7). □

Similar to the proof of above theorem, simple computations give the next theorem.

THEOREM 6. *If $g \in C_B[0,\infty)$ and $x \in [0,\infty)$, we get*

$$|T_n(g;x) - g(x)| \leq M\omega_2\left(g; \frac{1}{2}\sqrt{\frac{1}{n^2}x\left(8x^3\alpha^2 + 4x\alpha + n\right) + \frac{2\mu x}{n}\frac{e_\mu(-nx)}{e_\mu(nx)}}\right) + \omega\left(g; \frac{2\alpha x^2}{n}\right)$$

where M is a positive constant.

REMARK 1. *The case of $\mu = 0$ in Theorem 6 gives the result given in* [**14**].

References

[1] Altın, A., Doğru, O., Taşdelen, F. The generalization of Meyer-König and Zeller operators by generating functions, J. Math. Anal. Appl., 312 (1) **(2005)**, 181-194.

[2] Altomare, F., Campiti, M. Korovkin-Type Approximation Theory and its Applications, de Gruyter Studies in Mathematics, vol. 17, Walter de Gruyter, Berlin, Germany, **1994**.

[3] Appell, P., Kampe de Feriet, J. Hypergeometriques et Hyperspheriques: Polynomes d'Hermite, Gauthier-Villars, Paris, 1926.

[4] Atakut, Ç., İspir, N. Approximation by modified Szász–Mirakjan operators on weighted spaces, Proc. Indian Acad. Sci. Math. 112 **(2002)**, 571–578

[5] Atakut, Ç., Büyükyazici, İ. Stancu type generalization of the Favard Szász operators, Appl. Math. Lett., 23 (12) **(2010)**, 1479-1482.

[6] Ciupa, A. A class of integral Favard–Szász type operators. Stud. Univ. Babes-Bolyai Math. 40 (1) **(1995)**, 39–47.

[7] DeVore, R.A., Lorentz, G.G. Construtive Approximation, Springer, Berlin, **1993**.

[8] Doğru, O., Özarslan, M.A., Taşdelen, F. On positive operators involving a certain class of generating functions, Studia Sci. Math. Hungar., 41 (4) **(2004)**, 415-429.

[9] Gadzhiev, A.D. The convergence problem for a sequence of positive linear operators on unbounded sets and theorems analogues to that of P.P. Korovkin, Sov. Math. Dokl. 15 (5) **(1974)**, 1453-1436.

[10] Gupta, V., Vasishtha, V., Gupta, M.K. Rate of convergence of the Szász–Kantorovich–Bezier operators for bounded variation functions, Publ. Inst. Math. (Beograd) (N.S.), 72 **(2006)**, 137–143.

[11] İcoz, G., Cekim, B. Dunkl generalization of Szász operators via q-calculus, Journal of Inequalities and Applications, 2015:284 **(2015)**, 11 pages.

[12] İcoz, G., Cekim, B. Stancu-type generalization of Dunkl analogue of Szász–Kantorovich operators, Math. Meth. Appl. Sci, 39 **(2016)**, 1803–1810.

[13] Korovkin, P. P. On convergence of linear positive operators in the space of continuous functions (Russian), Doklady Akad. Nauk. SSSR (NS) 90 **(1953)**, 961–964.

[14] Krech, G. A note on some positive linear operators associated with the Hermite polynomials, Carpathian J. Math., 32 (1) **(2016)**, 71–77.

[15] Mursaleen, M., Rahman, S., Alotaibi. A. Dunkl generalization of q-Szász-Mirakjan Kantorovich operators which preserve some test functions, Journal of Inequalities and Applications, 2016:317 **(2016)**, 18 pages.

[16] Mursaleen, M., Nasiruzzaman, Md., Alotaibi. A.On modified Dunkl generalization of Szász operators via q-calculus, Journal of Inequalities and Applications, 2017:38 **(2017),** 12 pages.

[17] Olgun, Ali, İnce, H. Gül, Taşdelen, Fatma. Kantrovich type generalization of Meyer-König and Zeller operators via generating functions, An. Ştiinţ. Univ. "Ovidius" Constanţa Ser. Mat., 21 (3) **(2013)**, 209–221.

[18] Rosenblum, M. Generalized Hermite polynomials and the Bose-like oscillator calculus, Oper. Theory: Adv. Appl. 73 **(1994)**, 369-396.

[19] Stancu, D.D. Approximation of function by a new class of polynomial operators, Rev. Rourn. Math. Pures et Appl., 13 (8) **(1968)**, 1173-1194.

[20] Sucu, S. Dunkl analogue of Szász operators, Appl. Math. Comput. 244 **(2014)**, 42-48.

[21] Sucu, S., İçöz, G., Varma, S. On some extensions of Szasz operators including Boas-Buck type polynomials, Abstr. Appl. Anal., Vol. **2012**, Article ID 680340, 15 pages.

[22] Szász, O. Generalization of S. Bernstein's polynomials to the infinite interval, J. Res. Nat. Bur. Stand. 45 **(1950)**, 239–245.

[23] Taşdelen, F., Aktaş, R., Altın,A. A Kantrovich Type of Szasz Operators Including Brenke-Type Polynomials, Abstract and Applied Analysis, Vol. 2012 **(2012)**, 13 pages.
[24] Varma, S., Sucu, S., İçöz, G. Generalization of Szasz operators involving Brenke type polynomials, Comput. Math. Appl., 64 (2) **(2012)**, 121-127.
[25] Varma, S., Tasdelen F. Szasz type operators involving Charlier polynomials, Mathematical and Computer Modeling, 56 **(2012)**, 118-122.

Ankara University, Faculty of Science, Department of Mathematics, 06100, Tandoğan, Ankara, Turkey
E-mail address: raktas@science.ankara.edu.tr

Current address: Gazi University, Faculty of Science, Department of Mathematics, 06100, Beşevler, Ankara, Turkey
E-mail address: bayramcekim@gazi.edu.tr

Ankara University, Faculty of Science, Department of Mathematics, 06100, Tandoğan, Ankara, Turkey
E-mail address: tasdelen@science.ankara.edu.tr